

## Finite Double Integral Representation for the Polynomial Set $T_n(X_1, X_2, X_3, X_4)$

Ahmad Quaisar Subhani

*Dept. of Maths, Maulana Azad College of Engg and technology, Anisabad, Patna*

**ABSTRACT:-** Recently, we introduced “An unification of certain generalized Geometric polynomial Set  $T_n(x_1, x_2, x_3, x_4)$ , with the help of generating function which contains Appell function of four variables in the notation of Burchall and Chaundy [4] associated with Lauricella function. This generated hypergeometric polynomial Set covers as many as thirty four orthogonal and non-orthogonal polynomials. In the present paper an attempt has been made to express a double finite integral representation of the polynomial set  $T_n(x_1, x_2, x_3, x_4)$ .

**AMS Subject Classification:-** Special function-33

**Keywords:-** Appell function, Lauricella form, Generalized hypergeometric polynomial.

### I. INTRODUCTION

The generalized polynomial set  $T_n(x_1, x_2, x_3, x_4)$  is defined by means of generating relation

$$e^{\lambda_1 t^{m_1} + d_\mu} F \left[ \begin{matrix} (A_r); (C_u); (E_{g_i}) \\ \lambda_1 t, \lambda_2 x_2^{m_2} t^{m_2}, \lambda_3 x_3^{m_3} t^{m_3}, \lambda_4 x_4^{m_4} t^{m_4} \\ (B_s); (D_v); (F_{q_i}) \end{matrix} \right] = \sum_{n=0}^{\infty} T_{n,m:m_1:m_2:m_3:m_4} (x_1, x_2, x_3, x_4) t^n \dots \quad (1.1)$$

where ( $i = 1, 2, 3, 4$ ) and  $\lambda, \lambda_1, \lambda_2, \lambda_3, \lambda_4$  are real and  $m, m_1, m_2, m_3, m_4$  are positive integer. The left hand side of (1.1) contains the product of two generalized hypergeometric function which contains Appell function of four variables in the notation of Burchall and Chaundy [4] associated with Lauricella function. The polynomial set contains a number of parameters for simplicity, it is denoted by  $T_n(x_1, x_2, x_3, x_4)$ , where  $n$  is the order of the polynomial set.

After little simplification (1.1) gives,

$$T_n(x_1, x_2, x_3, x_4) = \sum_{P_1=0}^{\left[\frac{n}{m_1}\right]} \sum_{P_2=0}^{\left[\frac{n-m_1 P_1}{m_2}\right]} \sum_{P_3=0}^{\left[\frac{n-m_1 P_1 - m_2 P_2}{m_3}\right]} \sum_{P_4=0}^{\left[\frac{n-m_1 P_1 - m_2 P_2 - m_3 P_3}{m_4}\right]} \times \frac{[(A_r)]_{n-m_1 P_1 - (m_2-1) P_2 - (m_3-1) P_3 - (m_4-1) P_4}}{[(B_s)]_{n-m_1 P_1 - (m_2-1) P_2 - (m_3-1) P_3 - (m_4-1) P_4}} \times \frac{[(C_u)]_{n-m_1 P_1 - m_2 P_2 - m_3 P_3 - m_4 P_4} [(E_{g_1})]_{P_1} [(E_{g_2})]_{P_2} [(E_{g_3})]_{P_3}}{[(D_v)]_{n-m_1 P_1 - m_2 P_2 - m_3 P_3 - m_4 P_4} [(F_{q_1})]_{P_1} [(F_{q_2})]_{P_2} [(F_{q_3})]_{P_3}} \frac{[(E_{g_4})]_{P_4} (\lambda x_1^m)^{n-m_1 P_1 - m_2 P_2 - m_3 P_3 - m_4 P_4} \lambda_1^{P_1} (\alpha_\mu)^{P_1} \lambda_2^{P_2} (\lambda_3 x_3^{m_3})^{P_3} (\lambda_4 x_4^{m_4})^{P_4}}{[(F_{q_4})]_{P_4} P_1! P_2! x_2^{m_2 P_2} P_3! P_4!} \dots \dots \quad (1.2)$$

The polynomial Set  $T_n(x_1, x_2, x_3, x_4)$  happens to be the generalization of as many thirty four orthogonal and non-orthogonal polynomials.

**Notations:**

a)

- I.  $(m)=1, 2, 3, \dots, m$
- II.  $(A_p) = A_1, A_2, A_3, \dots, A_p$
- III.  $[(A_p)] = A_1, A_2, A_3, \dots, A_p$
- IV.  $[(A_p)]_n = (A_1)_n, (A_2)_n, (A_3)_n, \dots, (A_p)_n$
- V.  $\Delta(a, b) = \frac{b}{a}, \frac{b+1}{a}, \dots, \frac{b+a-1}{a}$

b)

$$\text{I. } M = \frac{[(A_r)]_n [(C_u)]_n (\lambda x^m)^n}{[(B_s)]_n [(D_v)]_n n!}$$

**2. Theorem :** For  $m_2 > 1$ ,  $m_3 > 1$  and  $m_4 > 1$ , we get

$$T_n(x_1, x_2, x_3, x_4) = \frac{\Gamma(\mu + \nu) M_1}{\Gamma \mu \Gamma \nu} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} \eta^\mu (1-\eta)^{\nu-1}}{(1-\xi\eta)^{\mu+\nu-1}}$$

$$\times F_{r+u:q_1:q_2:q_3:q_4}^{1+s+\nu: g_1: g_2: g_3: g_4} \left[ \begin{matrix} [(-n): m_1, m_2, m_3, m_4] \\ \hline \end{matrix} \right]$$

$$[(1-(B_s)-n): m_1, m_2-1, m_3-1, m_4-1], [(1-(D_v)-n): m_1, m_2, m_3, m_4]$$

$$[(1-(A_r)-n): m_1, m_2-1, m_3-1, m_4-1], [(1-(C_u)-n): m_1, m_2, m_3, m_4]$$

$$[(E_{g_1}):1], [(E_{g_2}):1], [(E_{g_3}):1], [(E_{g_4}):1], \left[ \left( \frac{\mu+\nu}{2} \right):1 \right] \left[ \left( \frac{\mu+\nu+1}{2} \right):1 \right]$$

$$[(F_{q_1}):1], [(F_{q_2}):1], [(F_{q_3}):1], [(F_{q_4}):1], [(\mu):1], [(\nu):1]$$

$$\frac{\lambda_1 \alpha_\mu (-1)^{m_1(1+s+u+\nu+1)}}{(\lambda x_1^m)^{m_1}}, \frac{\lambda_2 (-1)^{\{m_2(r+s+u+\nu+1)+r+s\}}}{(\lambda x_1^m x_2)^{m_2}},$$

$$\frac{\lambda_3 x_3^{m_3} (-1)^{\{m_3(r+s+u+\nu+1)+r+s\}}}{(\lambda x_1^m)^{m_3}}, \frac{\lambda_4 x_4^{m_4} (-1)^{\{m_4(r+s+u+\nu+1)+r+s\}}}{(\lambda x_1^m)^{m_4}} \Bigg] d\xi d\eta$$

.... (2.1)

**Proof :** We have

$$\int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \times \sum_{P_1=0}^{\left[ \frac{n}{m_1} \right]} \sum_{P_2=0}^{\left[ \frac{n-m_1 P_1}{m_2} \right]} \sum_{P_3=0}^{\left[ \frac{n-m_1 P_1 - m_2 P_2}{m_3} \right]}$$

$$\begin{aligned}
 & \sum_{P_4=0}^{\left[ \frac{n-m_1P_1-m_2P_2-m_3P_3}{m_4} \right]} \frac{[(A_r)]_{n-m_1P_1-(m_2-1)P_2-(m_3-1)P_3-(m_4-1)P_4}}{[(B_s)]_{n-m_1P_1-(m_2-1)P_2-(m_3-1)P_3-(m_4-1)P_4}} \\
 & \times \frac{[(C_u)]_{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} \left[ (E_{g_1}) \right]_{P_1} \left[ (E_{g_2}) \right]_{P_2} \left[ (E_{g_3}) \right]_{P_3}}{[(D_v)]_{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} \left[ (F_{q_1}) \right]_{P_1} \left[ (F_{q_2}) \right]_{P_2} \left[ (F_{q_3}) \right]_{P_3}} \\
 & \times \frac{\left[ (E_{g_4}) \right]_{P_4} (\lambda x_1)^{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} (\lambda_1 \alpha_\mu)^{P_1} \lambda_2 P_2 (\lambda_3 x_3^{m_3})^{P_3}}{\left[ (F_{q_4}) \right]_{P_4} (n-m_1P_1-m_2P_2-m_3P_3-m_4P_4)! P_1 x_2^{m_2 P_2} P_2! P_3!} \\
 & \times \frac{(\lambda_4 x_4^{m_4})^{P_4} \left( \frac{\mu+\nu}{2} \right)_{P_1} \left( \frac{\mu+\nu+1}{2} \right)_{P_1} d\xi d\eta}{P_4! (\nu)_{P_1} (\mu)_{P_1}} \\
 & = \sum_{P_1=0}^{\left[ \frac{n}{m_1} \right]} \sum_{P_2=0}^{\left[ \frac{n-m_1P_1}{m_2} \right]} \sum_{P_3=0}^{\left[ \frac{n-m_1P_1-m_2P_2}{m_3} \right]} \sum_{P_4=0}^{\left[ \frac{n-m_1P_1-m_2P_2-m_3P_3}{m_4} \right]} \\
 & \times \frac{[(A_r)]_{n-m_1P_1-(m_2-1)P_2-(m_3-1)P_3-(m_4-1)P_4}}{[(B_s)]_{n-m_1P_1-(m_2-1)P_2-(m_3-1)P_3-(m_4-1)P_4}} \\
 & \times \frac{[(C_u)]_{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} \left[ (E_{g_1}) \right]_{P_1} \left[ (E_{g_2}) \right]_{P_2} \left[ (E_{g_3}) \right]_{P_3}}{[(D_v)]_{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} \left[ (F_{q_1}) \right]_{P_1} \left[ (F_{q_2}) \right]_{P_2} \left[ (F_{q_3}) \right]_{P_3}} \\
 & \times \frac{\left[ (E_{q_4}) \right]_{P_4} (\lambda_1 \alpha_\mu)^{P_1} \lambda_2^{P_2} (\lambda_3 x_3^{m_3})^{P_3} (\lambda_4 x_4^{m_4})^{P_4}}{\left[ (F_{q_4}) \right]_{P_4} P_1! P_2! x_2^{m_2 P_2} P_3! P_4!} \\
 & \times \frac{(\lambda x_1)^{n-m_1P_1-m_2P_2-m_3P_3-m_4P_4} \left( \frac{\mu+\nu}{2} \right)_{P_1} \left( \frac{\mu+\nu+1}{2} \right)_{P_2}}{(n-m_1P_1-m_2P_2-m_3P_3-m_4P_4)! (\nu)_{P_1} (\mu)_{P_1}} \\
 & \times \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1+P_1} (1-\eta)^{\nu-1+P_1} \eta^{\mu+P_1}}{(1-\xi\eta)^{\mu+\nu-1+2P_1}} d\xi dn \\
 & = \frac{\Gamma(\mu) \Gamma(\nu) M_1}{\Gamma(\mu+\nu)} \sum_{P_1, P_2, P_3, P_4=0}^{\infty} \frac{[1-(B_s)-n]_{m_1P_1+(m_2-1)P_2+(m_3-1)P_3+(m_4-1)P_4}}{[1-(A_r)-n]_{m_1P_1+(m_2-1)P_2+(m_3-1)P_3+(m_4-1)P_4}}
 \end{aligned}$$

$$\begin{aligned}
 & \times \frac{[1-(D_v)-n]_{m_1 P_1 + m_2 P_2 + m_3 P_3 + m_4 P_4} \left[ (E_{g_1}) \right]_{P_1} \left[ (E_{g_2}) \right]_{P_2}}{[1-(C_u)-n]_{m_1 P_1 + m_2 P_2 + m_3 P_3 + m_4 P_4} \left[ (F_{q_1}) \right]_{P_1} \left[ (F_{q_2}) \right]_{P_2}} \\
 & \times \frac{\left[ (E_{g_3}) \right]_{P_3} \left[ (E_{g_4}) \right]_{P_4} \left( \frac{\mu+v}{2} \right)_{P_1} \left( \frac{\mu+v+1}{2} \right)_{P_1} (\lambda_1 \alpha_\mu)^{P_1}}{\left[ (F_{q_3}) \right]_{P_3} \left[ (F_{q_4}) \right]_{P_4} (v)_{P_1} (\mu)_{P_1} P_1!} \\
 & \times \frac{(-1)^{m_1(r+s+u+v+1)P_1} \lambda_2^{P_2} (-1)^{\{m_2(r+s+u+v+1)+r+s\}P_2}}{(\lambda x_1^m)^{m_1 P_1} (\lambda x_1^m x_2)^{m_2 P_2} P_2!} \\
 & \times \frac{\lambda_3 x_3^{m_3} (-1)^{\{m_3(r+s+u+v+1)+r+s\}P_3} \lambda_4 x_4^{m_4} (-1)^{\{m_4(r+s+u+v+1)+r+s\}P_4}}{(\lambda x_1^m)^{m_3 P_3} P_3! (\lambda x_1^m)^{m_4 P_4} P_4!} \\
 & \times (-n)_{m_1 P_1 + m_2 P_2 + m_3 P_3 + m_4 P_4} \dots \dots \quad (2.2)
 \end{aligned}$$

The single terminating factor makes all summation in (2.2) runs up to  $\infty$ .

$$= \frac{\Gamma(\mu)}{\Gamma(\mu+v)} T_n(x_1, x_2, x_3, x_4)$$

On using [7],

$$\int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{v-1} \eta^\mu}{(1-\xi\eta)^{\mu+v-1}} d\xi d\eta = B(\mu:v)$$

#### Particular Cases :

(i) On taking  $r=0=s=u=v=g_3$ ;  $q_3=1=x_3=m=\lambda=\lambda_2=\alpha_\mu$ ;  $F_1=\lambda+\frac{1}{2}$ ;  $m_3=2$

and writing  $\frac{x}{\sqrt{x^2-1}}$  for  $x$ , in (2.1), we have

$$\begin{aligned}
 C_n^\lambda(x) &= \frac{(2\lambda)_\eta \left( x^2 - 1 \right)^{-n/2} x^n \Gamma(\mu+v)}{n! \Gamma\mu \Gamma v} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} \eta^\mu (1-\eta)^{v-1}}{(1-\xi\eta)^{\mu+v-1}} \\
 &\quad \times {}_2F_1 \left[ \begin{matrix} -\frac{n}{2}, -\frac{n}{2} + \frac{1}{2}, \frac{\mu+v}{2} \\ \frac{\mu+v+1}{2}; \quad \frac{x^2-1}{x^2} \\ \alpha + \frac{1}{2}, \mu, v; \end{matrix} \right] d\xi d\eta
 \end{aligned}$$

(ii) If we take  $s = 0 = u = v = g_3 = q_3; m = 1 = r = x_3 = \lambda_2 = \alpha_\mu; \lambda = 2; m_3 = 3;$

$A_1 = \frac{1}{2}$  and  $x$  for  $x_1$ , in (2.1), we have

$$g_n^*(x) = \frac{\left(\frac{1}{2}\right)_n 3^n x^n \Gamma(\mu + \nu)}{n! \Gamma(\mu) \Gamma(\nu)} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times F \left[ \begin{array}{c} \Delta(3; m), \frac{\mu+\nu}{2} \\ \frac{\mu+\nu+1}{2}; \quad \frac{1}{4x^3} \\ \Delta\left(2; \frac{1}{2}-n\right), \nu, \mu; \end{array} \right] d\xi dn$$

(iii) On putting  $r = 0 = s = u = v = q_4 = g_4; x_4 = 1 = m = \alpha_\mu; \lambda_3 = h;$

$\lambda = m^{-m}, x_1 = x$  in (2.1), we obtain

$$g_n^m(x, h) = \frac{\Gamma(\mu + \nu) x^n}{\Gamma(\mu) \Gamma(\nu) n!} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times F \left[ \begin{array}{c} \Delta(m; -n), \frac{\mu+\nu}{2}; \\ \frac{\mu+\nu+1}{2}; \quad h\left(\frac{m}{2}\right)^m \\ \mu, \nu; \end{array} \right] d\xi d\eta$$

(iv) On making the substitution  $s = 0 = u = v = g_4 = q_4; m = 1 = r = \alpha_\mu, A_1 = -P;$

$m_4 = m; x_4 = y^{\frac{1}{m}}, \lambda = \{-(m-1)\}^{\frac{m-1}{m}}, \lambda_4 = (-1)^{m+1} c^{m-1}$  and  $x_1 = x$  in (2.1), we obtain

$$P_n(m, x, y, p, c) = \frac{\Gamma(\mu + \nu) \{-(m-1)\}^{\frac{n(m-1)}{m}} (mn)^n C^{Pn}(-p)_n}{\Gamma(\mu) \Gamma(\nu) n!} \\ \times \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times F \left[ \begin{array}{c} \Delta(m; -n); \frac{\mu+\nu}{2}, \frac{\mu+\nu+1}{2} \\ \Delta(m-1; 1-p-n); \mu, \nu; \end{array} \frac{(-1)^{m+1} C^{m-1} y}{x^m \{-(m-1)\}^{m-1}} \right] d\xi dn$$

(v) On taking  $r = 0 = s = u = v = g_4 = q_4; x_4 = 1 = m = \lambda = \alpha_\mu; m_4 = P; \lambda_g = -1$  and  $x_1 = x$  in (2.1), we get

$$g_n^P(x) = \frac{\Gamma(\mu + \nu)}{\Gamma(\mu)} \frac{p^n}{\Gamma(\nu)} \frac{x^n}{n!} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times F \left[ \begin{matrix} \Delta(p; -n), \frac{\mu + \nu}{2}, \frac{\mu + \nu + 1}{2}; & -\left(-\frac{1}{x}\right)^P \\ \mu, \nu; & \end{matrix} \right] d\xi d\eta$$

(vi) If we get  $r = 0 = s = u = v = g_4 = q_4 = 1 = m = \alpha_\mu = x_4; \lambda = \frac{\nu}{m}; \lambda_4 = \mu$ ;

$m_4 = m; x_1 = x$  and replacing  $E_1 = (a_r)$  and  $F_1 = (b_s)$  in (2.1), we get

$$A_n(x) = \frac{\Gamma(\mu + \nu)}{\Gamma(\mu)} \frac{(\nu x)^n}{\Gamma(\nu)} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times F \left[ \begin{matrix} \Delta(m; -n), (a_r), \frac{\mu + \nu}{2}, \frac{\mu + \nu + 1}{2}; & \mu \left(\frac{-m}{\nu x}\right)^m \\ (bs), \mu, \nu; & \end{matrix} \right] d\xi d\eta$$

(vii) On taking  $r = 0 = s = u = v; e_4 = P; m = -\frac{1}{p}; \lambda = 1 = \lambda_4 = \alpha_\mu; x_4 = -1$  and

$x_1 = x$  and writing  $E_{g_4} = \alpha_u, F_{q_4} = \beta_v$  in (2.1), we get

$$B_n^P(x) = \frac{\Gamma(\mu + \nu)}{\Gamma(\mu)} \frac{(-p\alpha x^y)^n}{\Gamma(\nu)} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times {}_{P+u}F_v \left[ \begin{matrix} \Delta(p; -n), (\alpha_u) \\ \frac{\mu + \nu}{2}, \frac{\mu + \nu + 1}{2}; x \end{matrix} \right] d\xi d\eta \\ \left[ (\beta v), \mu, \nu; \right]$$

(viii) On making the Substitution  $r = 0 = s = u = v = g_4 = q_4$ ;

$x_1 = 1 = m = \alpha_\mu; \lambda_4 = \beta, x_a = m = m_4$  and  $\lambda = \alpha$  in (2.1), we get

$$H_{n,m}(\alpha, \beta) = \frac{\Gamma(\mu + \nu) \cdot \alpha^\eta}{\Gamma(\mu) \Gamma(\nu)} \int_0^1 \int_0^1 \frac{(1-\xi)^{\mu-1} (1-\eta)^{\nu-1} \eta^\mu}{(1-\xi\eta)^{\mu+\nu-1}} \\ \times {}_mF_0 \left[ \begin{matrix} \Delta(m; -n), \frac{\mu + \nu}{2}, \frac{\mu + \nu + 1}{2} & \frac{\beta(-m)^m}{\alpha} \\ \mu, \nu; & \end{matrix} \right] d\xi d\eta$$

#### **REFERENCES**

- [1]. Agarwal, R.P. (1965) An extension of Meijer's G-function, Proc. Nat. Institute Sci. India, 31, a, 536-546.
- [2]. Abdul Halim, N and A characterization of Laguerre polynomials, Al\_Salam, W.A. (1964) Rend, Sem, Univ., padova, 34,176 -179.
- [3]. Agarwal, B.D. and On a multivariate polynomials system Mukherjee, S. (1988) proceedings of national symposium on special functions and their applications. Gorakhpur (india) p. 153 -158.
- [4]. Burchnall, J.L. and Expansions of Appell's double hyper Chaundy, T.W. (1941) geometric functions (ii), Quart. J. Math. Oxford ser. 12,112 -128
- [5]. Carlitz, L. and Some hyper geometric polynomials Shrivastava, H.M. (1976) associated with the Lauricella function ED of several variables
- [6]. Singh, Ramji and Unification of classical polynomials. Math. Singh, D.N. (1982) Edu. Vol. 16, no. 3. Sept. p. 69 - 77.
- [7]. Edwards, J. (1964) Treatise on integral Calculus, vols. I, II, Chelsea Publishing Company, New York.